



中国科学技术大学
University of Science and Technology of China

GAMES 301：第10讲

共形参数化1

Spin变换、Circle填充 & 共轭调和函数

方清
中国科学技术大学

Content



- 1. Introduction to conformal mapping**
- 2. Differential of conformal mapping**
- 3. Spin transformations**
- 4. Circle packing and circle patterns**
- 5. Conjugate harmonic functions**

1

Introduction to conformal mapping

創寰宇學府
育天下英才
原濟昌月

Angle-based flattening (ABF)



- Key observation: the parameterized triangles are uniquely defined by all the angles at the corners of the triangles.
 - Find angles instead of uv coordinates.
 - Use angles to reconstruct uv coordinates.


- Angel preservation:

- Interior vertex:

$$\beta_i^{jk} = \frac{\alpha_i^{jk} \cdot 2\pi}{\sum_i \alpha_i^{jk}}.$$

- Boundary vertex:

$$\beta_i^{jk} = \alpha_i^{jk}.$$



Angle-based flattening (ABF)




- Least square optimization:

$$\min_{\theta > 0} \sum_{ijk} (\beta_i^{jk} - \theta_i^{jk})^2 + (\beta_j^{ki} - \theta_j^{ki})^2 + (\beta_k^{ij} - \theta_k^{kj})^2$$

s.t.

$$\left\{ \begin{array}{l} \sum_{t_{ijk} \in St(i)} \theta_i^{jk} = 2\pi, \quad \forall i \text{ interior vertex} \\ \\ \theta_i^{jk} + \theta_j^{ki} + \theta_k^{ij} = \pi, \quad \forall t_{ijk} \\ \\ \prod_{t_{ijk} \in St(i)} \frac{\sin \theta_j^{ki}}{\sin \theta_k^{ij}} = 1, \quad \forall i \text{ interior vertex} \end{array} \right.$$




$$\frac{\sin \theta_j^{ki}}{\sin \theta_k^{ij}} = \frac{l_{ki}}{l_{ij}}$$

Least-square conformal mapping (LSCM)



- Angle preservation → similar transforms



Least-square conformal mapping (LSCM)



- Angle preservation → similar transform



- For triangle t_{ijk} :

$$\begin{pmatrix} du \\ dv \end{pmatrix} = \begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix}$$



$$s \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} M$$

$$\Rightarrow \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$






Least-square conformal mapping (LSCM)



- Least square optimization:

$$E_{LSCM} = \sum_{ijk} A_{ijk} \left(\left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right)^2 + \left(\frac{\partial u}{\partial y} + \frac{\partial v}{\partial x} \right)^2 \right)$$



- Not unique minimizer → fixing at least two vertices.



Conformal mapping






- Angle preservation



Applications




- Texturing
- Morphing
- Remeshing
- Shape analysis
- ...



Applications





- Cartography
- Architecture
- Art design
- Fabrication
- ...



Applications



- Fluids
- Microstructures
- Topology optimization
- ...






Differential of conformal mapping

創寰宇學府
育天下英才
原濟昌月


Conformal mapping



- Discrete conformal mapping.
 - Angle of triangles.
 - Piecewise linear function.



$$\frac{l_{jk}}{\sin \theta_i^{jk}} = \frac{l_{ki}}{\sin \theta_j^{ki}} = \frac{l_{ij}}{\sin \theta_k^{ij}} = 2R$$





“Rigid”

Conformal mapping




- Discrete conformal mapping.
 - Angle of triangles.
 - Piecewise linear function.
- Continuous conformal mapping.
 - Angle between vectors.
 - Smooth function.



Differential



- Push forward of vector: $df(v) = \lim_{t \rightarrow 0} \frac{f(p+tv)-f(p)}{t}$
- Linear operator: $df(v) = J_f v$, J_f Jacobian matrix.





Differential



- Angle preservation:
 - $\theta[v, w] \Leftrightarrow \theta[df(v), df(w)]$

- Complex number
 - Rotate & scale : $z = |z|e^{i\phi}$
 - $v \rightarrow w \Leftrightarrow w = vz = |v||z|e^{i(\theta+\phi)}$



$$\text{Arg}(v^{-1}w) = \text{Arg}(df^{-1}(v)df(w))$$

$$v^{-1}w = df^{-1}(v)df(w)$$

$$\text{Complex linear: } df(zv) = zd(df(v))$$


Differential



- Let $f = f_x + i f_y$.

$$- df(i) = \lim_{t \rightarrow 0} \frac{f(x+yi+ti) - f(x+yi)}{t} = \frac{\partial f_x}{\partial y} + \frac{\partial f_y}{\partial y} i$$


$$- df(1) = \lim_{t \rightarrow 0} \frac{f(x+yi+t) - f(x+yi)}{t} = \frac{\partial f_x}{\partial x} + \frac{\partial f_y}{\partial x} i$$



- Cauchy-Riemann equation: $df(i) = i df(1)$

$$- \begin{cases} \frac{\partial f_x}{\partial x} = \frac{\partial f_y}{\partial y} \\ \frac{\partial f_x}{\partial y} = -\frac{\partial f_y}{\partial x} \end{cases} \Leftrightarrow J_f = \begin{pmatrix} a & -b \\ b & a \end{pmatrix}$$

$$- \forall v \in \mathbb{C}, df(v) = df(1)v$$




Similar transform!

Differential




- Plane to plane : $df(v) = \lim_{t \rightarrow 0} \frac{f(p+tv)-f(p)}{t}$
- Manifold to manifold
 - Curve : $\Gamma(0) = p, \Gamma'(0) = v$
 - $df(v) = \lim_{t \rightarrow 0} \frac{f(\Gamma(t))-f(\Gamma(0))}{t} \quad \rightarrow \quad df(v) : v \in T_p M \rightarrow w \in T_{f(p)} N$



Differential



- Cauchy-Riemann equation
 - Plane : $df(i) = idf(1)$
 - Manifold : $df(J_M v) = J_N df(v), \forall v \in T_p M$



3



Spin transformation

創寰宇學府
育天下英才
原濟昌月
一九八六年五月

Quaternions



- Conformal mapping : rotate and scale
 - 2D plane : complex number
 - Surface embedding in \mathbb{R}^3 : quaternions






Quaternions

- From \mathbb{R}^3 to \mathbb{H} : $\vec{x} = (x_1, x_2, x_3) \rightarrow x = (0, \vec{x}) = 0 + x_1\mathbf{i} + x_2\mathbf{j} + x_3\mathbf{k}$
- Rotation around the axis $\vec{u} = (u_1, u_2, u_3)$, $\|\vec{u}\| = 1$
 - $q = \left(\cos \frac{\theta}{2}, -\sin \frac{\theta}{2} \vec{u}\right) = \cos \frac{\theta}{2} - \sin \frac{\theta}{2} (u_1 \mathbf{i} + u_2 \mathbf{j} + u_3 \mathbf{k})$
 - $\bar{q} = \left(\cos \frac{\theta}{2}, \sin \frac{\theta}{2} \vec{u}\right)$, $q\bar{q} = 1 \rightarrow \bar{q} = q^{-1}$
 - $y = \bar{q}xq = 0 + y_1i + y_2j + y_3k \rightarrow \vec{y} = (y_1, y_2, y_3)$
- Scale
 - $q' = cq, c \in \mathbb{R} \Rightarrow y' = \bar{q}'xq = c^2y$

$$\mathbf{i}^2 = \mathbf{j}^2 = \mathbf{k}^2 = \mathbf{ijk} = -1$$



Spin transformation



- Spin equivalence:

- $f: M \rightarrow \mathbb{R}^3$

- $\tilde{f}: M \rightarrow \mathbb{R}^3$

- $d\tilde{f}(X) = \bar{\lambda} df(X)\lambda, \exists \lambda: M \rightarrow \mathbb{H}$

- Dirac equation (integrable condition):


- $D\lambda = -\frac{df \wedge d\lambda}{|df|^2}$

- $(D - \rho)\lambda = 0, \exists \rho: M \rightarrow \mathbb{R}$

- Given initial ρ , solve λ (eigenvalue problem):

- $(D - \rho)\lambda = \gamma\lambda$



- $\rho \rightarrow \rho + \lambda$



Spin transformation



- Mean curvature half-density:
 - $(D - \rho)\lambda = 0, \exists \rho, \lambda: M \rightarrow \mathbb{R}$
 - $d\tilde{f}(X) = \bar{\lambda}df(X)\lambda \rightarrow \tilde{H}|d\tilde{f}| = H|df| + \rho|df|$
- Relation to conformal equivalence
 - Spin \rightarrow conformal
 - Conformal \nrightarrow spin




Circle packing and circle patterns

創寰宇學府
育天下英才
原濟昌月

Circle packing




- Smooth: infinitesimal circles preservation
- Discrete: preserve circles associated with mesh elements



Circle packing





- Limit → smooth conformal map



Circle packing



- For a triangulation K , $P = \{c_v\}$ is the circle packing of K if:
 - The center of $c_v \Leftrightarrow v \in V \subset K$
 - $\forall e_{ij} = v_i v_j$, c_{v_i}, c_{v_j} are tangent
 - $\forall f_{ijk} = v_i v_j v_k$, $c_{v_i}, c_{v_j}, c_{v_k}$ form a positively oriented triple




Circle packing



- Necessary and Sufficient Condition:

Given a triangulation K of a topological disk and a constraint radius at each boundary vertex, there is an (essentially) unique circle packing realizing the boundary constraints, with interior angles summing to 2π .

$$\cos \theta_t = \frac{(r_0 + r_1)^2 + (r_0 + r_2)^2 - (r_1 + r_2)^2}{2(r_0 + r_1)(r_0 + r_2)}$$




Circle packing



- Algorithm: repeat

For each $v_i \in V^\circ$:


- Let θ be total angle currently covered by k neighbors
 - Let r be radius such that k neighbors of radius r also cover θ
 - Set new radius of c_{v_i} such that k neighbors of radius r cover 2π
-



Circle packing





- Lack of geometry information



Circle patterns



- Associate each face with its circumcircle
- Preserve circle intersection angles




Circle patterns



- For planar Delaunay triangulation


$$\forall e_{ij} \in E : \alpha_e = \begin{cases} \pi - \theta_k^{ij} - \theta_l^{ij}, & \text{for interior edges} \\ \pi - \theta_k^{ij}, & \text{for boundary edges} \end{cases}$$



Circle patterns



- For planar Delaunay triangulation
 - $\forall e_{ij} \in E : 0 < \alpha_e < \pi$
 - $\forall v_i$ interior vertices : $\sum_{e \ni v_i} \alpha_e = 2\pi$
 - $\forall v_i$ boundary vertices : $\sum_{e \ni v_i} \alpha_e = 2\pi - \kappa_i$



Circle patterns





- For planar Delaunay triangulation

- $\forall e_{ij} \in E : 0 < \alpha_e < \pi$

- $\forall v_i$ interior vertices : $\sum_{e \ni v_i} \alpha_e = 2\pi$

- $\forall v_i$ boundary vertices : $\sum_{e \ni v_i} \alpha_e = 2\pi - \kappa_i < 2\pi$

Local Delaunay!



Circle patterns




- Parameterization

- Input Delaunay triangulation :

$$\sum_{e \ni v_i} \alpha_e ! = 2\pi$$

- UV mesh :

$$\hat{\alpha}_e > 0 \text{ and } \sum_{e \ni v_i} \hat{\alpha}_e = 2\pi$$



- Coherent angle system for $\hat{\alpha}_e$:

$$\exists \hat{\theta}_k^{ij}, \text{ s.t.}$$

- $\hat{\theta}_k^{ij} > 0$

- $\forall t_{ijk} \in T, \hat{\theta}_k^{ij} + \hat{\theta}_i^{jk} + \hat{\theta}_j^{ki} = \pi$

- $\forall e_{ij} \in E :$

$$\hat{\alpha}_e = \begin{cases} \pi - \hat{\theta}_k^{ij} - \hat{\theta}_l^{ij}, & \text{for interior edges} \\ \pi - \hat{\theta}_k^{ij}, & \text{for boundary edges} \end{cases}$$

Optimize $\hat{\theta}_k^{ij}!$

$$\min_{\hat{\theta}_k^{ij}} \sum \left(\hat{\theta}_k^{ij} - \theta_k^{ij} \right)^2$$

5

Conjugate harmonic functions

創寰宇學府
育天下英才
原濟昌月

Conjugate harmonic functions



- Cauchy-Riemann equation on complex plane


- $$\begin{cases} \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \\ \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x} \end{cases} \Leftrightarrow \nabla v = i \nabla u$$

- Harmonic functions


- $$\begin{cases} \Delta u = \nabla \cdot (\nabla u) = \nabla \cdot (-i \nabla v) = 0 \\ \Delta v = \nabla \cdot (\nabla v) = \nabla \cdot (i \nabla u) = 0 \end{cases}$$

- Discretization on triangular meshes

- Edges (conjugate harmonic 1-forms)
- Vertices (conjugate harmonic coordinates)



1-form ω



uv coordinates

Conjugate harmonic 1-forms



- Harmonic 1-forms:

- $\forall t_{ijk}, \omega_{ij} + \omega_{jk} + \omega_{ki} = 0$
- $\forall v_i \text{ interior vertex}, \sum_{e_{ij} \ni v_i} \alpha_{ij} \omega_{ij} = 0$





- Dimension of harmonic 1-forms:

- Genus $g \Rightarrow \{\omega^{(1)}, \dots, \omega^{(2g)}\}$
- Homology basis P_1, \dots, P_{2g} :

$$\sum_{e_{ij} \in P_k} \omega_{ij} = c_k, k = 1, \dots, 2g$$

- Conjugate gradients

- $\{{}^* \omega^{(1)}, \dots, {}^* \omega^{(2g)}\}$
- Integrate $\omega^{(k)} + \sqrt{-1} {}^* \omega^{(k)}$



Conjugate harmonic coordinates



- Solving Laplacian equations:

- For interior vertices $\begin{cases} \Delta u = 0 \\ \Delta v = 0 \end{cases}$
- Boundary control

- Dirichlet boundary condition:


- Boundary curve $\gamma: \partial M \rightarrow \mathbb{R}^2$

$$u \Big|_{\partial M} = \gamma_u, v \Big|_{\partial M} = \gamma_v$$


- Neumann boundary condition:

- Boundary gradients $h: \partial M \rightarrow \mathbb{R}^2$

$$\partial_M u = h_u, \partial_M v = h_v$$



Harmonic, not conformal



Conformal, conjugate gradients



中国科学技术大学
University of Science and Technology of China

谢 谢 !

